



Feeding Your 4-H Meat Goat



What Is Your Goal?

- **County Fair**
- **State Fair**
- **KJLS**
- **National Show**

Facilities

Easily Accessible

Clean Bedding



Air Flow



Plenty of Space to Move Around

Shade



Shelter




Salt Block

Feeder (Height)



Clean Fresh Water



GOAT 14
For All Classes Of Goats


— GUARANTEED ANALYSIS —

CRUDE PROTEIN, minimum -----14.0%
(Including EQUIVALENT CRUDE PROTEIN
from NON-PROTEIN NITROGEN) -----1.3%
CRUDE FAT, minimum -----2.5%
CRUDE FIBER, maximum -----14.0%
PHOSPHORUS (P), minimum -----0.3%
CALCIUM (Ca), minimum -----0.4%, maximum-----1.0%
SALT (NaCl), minimum -----0.5%, maximum -----1.0%
COPPER (Cu), minimum -----20ppm
SELENIUM (Se), minimum -----0.25ppm
VITAMIN A, minimum -----1,800 I.U.-lb.

— INGREDIENTS —
GRAIN PRODUCTS, ROUGHAGE PRODUCTS, FORAGE PRODUCTS, PLANT PROTEIN PRODUCTS, MOLASSES PRODUCTS, AMMONIUM CHLORIDE, CALCIUM CARBONATE, SALT, MONO AND DICALCIUM PHOSPHATE, MAGNESIUM OXIDE, ZINC OXIDE, COPPER SULFATE, BASIC COPPER CHLORIDE, MANGANOUS OXIDE, IRON OXIDE, SODIUM SELENITE, ETHYLENEDIAMINE DIHYDRIDIDE, COBALT CARBONATE, THIAMINE MONONITRATE, VITAMIN A SUPPLEMENT, VITAMIN D-3 SUPPLEMENT, VITAMIN E SUPPLEMENT, NATURAL AND ARTIFICIAL FLAVORS.

— FEEDING DIRECTIONS —
Feed as the concentrate portion of the diet for growing and mature goats. Additional good quality hay or grazing should be provided.

APR



ADVANCER® PLUS MON20
MEDICATED
Show Goat Complete Feed

For the prevention of oesophagitis caused by *Eimeria cuniculi*, *Eimeria christersoni* and *Eimeria vivax* in goats maintained in confinement.

Monensin 20 G/TON

ACTIVE DRUG INGREDIENTS		G/TON	
Monensin	20	G/TON	

GUARANTEED ANALYSIS			
Crude Protein, min	18.0	%	
This includes not more than 1.4% equivalent protein from non-protein nitrogen.			
Crude Fat, min	2.5	%	Salt, max 1.4 %
Crude Fiber, max	12.0	%	Copper, min 25 PPM
Acid Detergent Fiber, max	10.0	%	Copper, max 55 PPM
Calcium, min	0.7	%	Selenium, min 0.3 PPM
Calcium, max	1.2	%	Vitamin A, min 8,000 I.U./lb
Phosphorus, min	0.3	%	Vitamin D, min 300 I.U./lb
Salt, min	0.5	%	Vitamin E, min 150 I.U./lb

INGREDIENTS
grain products, plant protein products, roughage products, processed grain by-products, forage products, molasses products, calcium carbonate, vegetable oil, urea, ammonium chloride, sodium sesquicarbonate, Acetivital, cryose fermentation extract, sodium niobate, calcium chloride, sodium calcium methylenediphosphate, potassium chloride, vitamin E supplement, L-lysine, natural flavor, artificial flavor, zinc oxide, potassium permanganate, sulfadiazine, sodium hydroxide, yeast, sodium dihydrogen phosphate, yeast, vitamin A supplement, manganese propionate, mineral oil, vitamin D-3 supplement, niacin supplement, calcium pantothenate, copper propionate, methacaine sodium lauryl ether complex (source of vitamin K activity), riboflavin supplement, cobalt propionate, ethylenediamine dihydrochloride, vitamin B-12 supplement, pyridoxine hydrochloride, cobalt carbonate, biotin, nicotinic acid, niacinamide, calcium nicotinate.

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance for copper.
This product has been formulated specifically for goats and is not intended for other species.

CAUTION: USE AS DIRECTED
Caution: Follow label directions. Feeding this feed containing added selenium at levels higher than recommended is not permitted.

FEEDING DIRECTIONS:
Feed Show Rite® Advancer® Plus MON20 continuously at the rate shown to non-lactating goats. A source of good quality hay and constant supply of clean fresh water is also required.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Do not feed unaltered. Monensin medicated cattle and goat feeds are safe for cattle and goats only. Consumption by unapproved species may result in toxic reactions. Must be thoroughly mixed in feed before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding unaltered or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not feed to lactating goats.

Hubbard does not use "Restricted-use Products" in their products and is in compliance with FDA and state requirements regarding the use, handling and storage of "Restricted-use Products".


HUBBARD FEEDS
111 West Cherry Street Suite 500
Mankato, MN 56001
Net Weight 50 lb (22.68 kg) or bulk invoice.

HUBBARD **47855**

REV 12/07

BO:1905730

8 48560 01359 0



- Pellet or Textured
- Complete (Balanced)
- Fresh
- READ the Label
 - Warnings?
 - Cautions?
 - Medicated
 - Non-Medicated
 - Ca - P (2:1)
- Roughage



Feed Storage



How Much Do You Feed?



Don't Think it! - Ink It!

MONTH:						
SUNDAY	MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY
H*			H			W
H*			H			W
H*			H			W
H*			H			W
H*			H			W

NE 16 doe, 4 ewe, 1 kid
NW 2 buck
SW 16 doe, 2 kid

Check Water

Goal
County Fair
7/29/2022

First of month
- Trim Feet
- deworm
- drench

FEED		
Pen 1	1	cup + Sup 2X Day
Pen 2	1	Heaping cup + Sup 2X Day
Pen 3	2	3/4 cup + Sup 2X Day
Pen 4	4	cup 2X Day

WEIGHTS	
Pen 1	55
Pen 2	42
Pen 3	/

Goodno Livestock



**Record Weight(s)
Regularly**

Parasite Control

- How Often
- What to Use





Exercise Program

- **Little Athletes**
- **Short/Quick Burst of Energy**
 - **Sprints vs Marathon**
 - **Driveway - Track - Treadmill**
 - **Good Flat Ground**
- **DON'T Over Do It!**

QUESTIONS?